ANALYTICA CHIMICA ACTA

An international journal devoted to all branches of analytical chemistry

(Full texts are incorporated in CJELSEVIER, a file in the Chemical Journals Online database available on STN International; Abstracted, indexed in: Aluminium Abstracts; Anal. Abstr.; Biol. Abstr.; BIOSIS; Chem. Abstr.; Curr. Contents Phys. Chem. Earth Sci.; Engineered Materials Abstracts; Excerpta Medica; Index Med.; Life Sci.; Mass Spectrum. Bull.; Material Business Alerts; Metals Abstracts; Sci. Citation Index)

Special Issue

1st International Cyber Congress on Analytical BioSciences, Japan, August 1997

VOL. 365 NOS. 1–3 CONTENTS June 5, 1998

Foreword 1

Microscopy

Simultaneous analysis of membrane potential and calcium mobilization in a pancreatic ß-cell line MIN6 by use of a double-probe imaging microscope-system
K. Ozawa, Y. Fukano, T. Azuma, Y. Masumoto, H. Hayashi, A. Tamura (Hiroshima, Japan), J. Miyazaki (Suita, Japan) and T. Masujima (Hiroshima, Japan) 3

Comparative analysis of topographic distribution of acetylated histone H4 by using confocal microscopy and a deconvolution system
T. Wako (Joetsu, Japan), M. Fukuda (Niigata, Japan), R. Furushima-Shimogawara (Tokyo, Japan), N.D. Belyaev, B.M. Turner (Birmingham, UK) and K. Fukui (Joetsu, Japan) 9

Nucleic acid structure

Atomic force microscopy of supercoiled DNA structure on mica
M. Tanigawa and T. Okada (Ibaraki, Japan) 19

Detection of higher-ordered DNA sequence by using terbium(III) luminescence
S. Sueda, T. Ihara, B. Juskowiak and M. Takagi (Fukuoka, Japan) 27

Highly sensitive assay of DNA abasic sites in mammalian cells-optimization of the aldehyde reactive probe method
A. Asaeda (Osaka, Japan), H. Ide, H. Terato (Higashi-Higashi, Japan), Y. Takamori and K. Kubo (Osaka, Japan) 35

Quantitative analysis by the polymerase chain reaction using fluorescence correlation spectroscopy
M. Kinjo (Sapporo, Japan) 43

Novel method of detecting mismatch on DNA by using photo-induced electron transfer through DNA
T. Ihara, J. Takata and M. Takagi (Fukuoka, Japan) 49

Prediction of an antisense-effective target site on VEGFmRNA by using RNase H and a library of random oligonucleotides
Y. Matsuda, K. Kamiya and K. Uchida (Ibaraki, Japan) 55

Molecular imprinting

Molecularly imprinted polymeric membranes involving tetrapeptide EQKL derivatives as chiral-recognition sites toward amino acids
M. Yoshikawa, T. Fujisawa, J. Izumi, T. Kitao and S. Sakamoto (Kyoto, Japan) 59

Chloramphenicol sensor based on an in-situ imprinted polymer
S. McNiven, M. Kato, R. Levi, K. Yano and I. Karube (Tokyo, Japan) 69

Synthesis of castasterone selective polymers prepared by molecular imprinting
A. Kugimiya, J. Matsui (Hiroshima, Japan), H. Abe (Tokyo, Japan), M. Aburatani (Toyama, Japan) and T. Takeuchi (Hiroshima, Japan) 75

(Continued overleaf)

CONTENTS DIRECT

ContentsDirect, the *free* e-mail alerting service, delivers the table of contents for this journal directly to your PC, prior to publication. The quickest way to register for **Contents Direct** is via the Internet at: **http://www.elsevier.nl/locate/ContentsDirect.** If you don't have access to the Internet you can register for this service by sending an e-mail message to cdsubs@elsevier.co.uk – specifying the title of the publication you wish to register for.

Molecular imprint membranes of polyacrylonitrile copolymers with different acrylic acid segments
T. Kobayashi, H. Ying Wang and N. Fujii (Niigata, Japan) 81
Molecular recognition in cinchona alkaloid molecular imprinted polymer rods
J. Matsui, I.A. Nicholls and T. Takeuchi (Hiroshima, Japan) 89

Other surface-binding techniques
In vitro selection of aptamers that recognize a monosaccharide
J. Kawakami, Y. Kawase and N. Sugimoto (Kobe, Japan) 95
Affinity adsorption separation of mutagenic molecules by polyacrylamide hydrogels comprising double-stranded DNA
D. Umeno, T. Kano and M. Maeda (Fukuoka, Japan) 101
Rapid method for assay of quantitative binding of soluble proteins and photosynthetic membrane proteins on poly(vinylidene difluoride) membranes
K. Aizawa and E. Gantt (College Park, MD, USA) 109
Direct observation of sample trapping process onto internal-surface reversed-phase pre-column
A. Tamura, M. Khademizadeh, K. Ozawa and T. Masujima (Hiroshima, Japan) 115

Immunoassays
Preparation of a specific monoclonal antibody against 2′-deoxycytidine
I. Darwish (Assiut, Egypt), T. Akizawa, K. Hirose, K. Omura (Osaka, Japan), N. EL-Rabbat (Assiut, Egypt) and M. Yoshioka (Osaka, Japan) 121
Preparation of monoclonal antibody against D-3-methoxy-4-hydroxyphenylglycol
K. Hirose, T. Akizawa and M. Yoshioka (Osaka, Japan) 129
Syntheses of antigens conjugated with 3-methoxy-4-hydroxyphenylglycol by Mannich reaction for enzyme immunoassay
K. Hirose, T. Akizawa, K. Asada, Y. Tanaka, Y. Negoro and M. Yoshioka (Osaka, Japan) 137
Monitor of antibodies in human saliva using a piezoelectric quartz crystal biosensor
I. Tajima, O. Asami and E. Sugiura (Aichi, Japan) 147
Enzyme-linked immunosorbent assay for plasma 24,25-dihydroxyvitamin D_3
T. Higashi, N. Kobayashi, H. Ohmi, Y. Hayashi and K. Shimada (Kanazawa, Japan) 151

Fluorescence techniques
Design, synthesis and characterization of a novel fluorescent probe for nitric oxide (nitrogen monoxide)
Y. Katayama, S. Takahashi and M. Maeda (Fukuoka, Japan) 159
Characterization of the fluorescence properties of 4-fluoro-7-nitrobenzo-2-oxa-1,3-diazole derivatives of some primary and secondary sympathomimetic amines
O. Al-Dirbashi, N. Kuroda and K. Nakashima (Nagasaki, Japan) 169
Fluorescence reaction of bilirubin with zinc ion in dimethyl sulfoxide and its application to assay of total bilirubin in serum
K. Kohashi, Y. Date, M. Morita and Y. Tsuruta (Hiroshima, Japan) 177
Formation of highly fluorescent adducts between 2′-deoxyguanosine and amino acids by ionizing radiation
H. Terato, M. Yoshimura, M. Hayashi, Y. Ohyama (Higashi-Hiroshima, Japan), O. Yamamoto (Hiroshima, Japan) and H. Ide (Higashi-Hiroshima, Japan) 183

Other irradiation techniques
Analysis of free-electron laser-induced infrared multiphoton dissociation for cholesterol ester
K. Awazu, A. Nagai, T. Tomimasu (Osaka, Japan) and K. Aizawa (Tokyo, Japan) 193
Light-independent isomerization of bacteriochlorophyll *g* to chlorophyll *a* catalyzed by weak acid in vitro
M. Kobayashi, T. Hamano, M. Akiyama (Tsukuba, Japan), T. Watanabe (Tokyo, Japan), K. Inoue (Hiratsuka, Japan), H. Oh-oka (Toyonaka, Japan), J. Amesz (Leiden, The Netherlands), M. Yamamura and H. Kise (Tsukuba, Japan) 199

Liquid chromatography with fluorescence detection
Determination of mevalonic acid in human urine by liquid chromatography with precolumn derivatization and fluorescence detection
S. Kawasaki, T. Nagaoka, Y. Saisho, A. Morimoto and T. Umeda (Osaka, Japan) 205
Liquid chromatographic determination of 5-methylcytosine in DNA with fluorescence detection
S. Sonoki, J. Lin and S. Hisamatsu (Kanagawa, Japan) 213
Simultaneous determination of serum and urinary hydroxyproline and proline by liquid chromatography using two fluorescent labeling reagents
H. Inoue, K. Kohashi and Y. Tsuruta (Hiroshima, Japan) 219
Serotonin monitoring in microdialysate from rat brain by microbore-liquid chromatography with fluorescence detection
J. Ishida (Fukuoka, Japan), T. Yoshitake, K. Fujino (Oita, Japan), K. Kawano (Fukuoka, Japan), J. Kehr (Stockholm, Sweden) and M. Yamaguchi (Fukuoka, Japan) 227
Simultaneous determination of 7-methylguanine and O^6-methylguanine in DNA by liquid chromatography with postcolumn fluorescence derivatization using 3,4-dimethoxyphenylglyoxal
Y. Ohba, M. Kai and K. Zaitsu (Fukuoka, Japan) 233

Chromatography and electrophoresis

Intelligent peptide-mapping method for recombinant human tissue-type plasminogen activator by lysyl-endopeptidase digestion
Y. Shikata, M. Kuwada, Y. Hayashi, A. Hashimoto, A. Koide and N. Asakawa (Ibaraki, Japan) 241

Separation of stereoisomers of C_{27}-bile acid CoA esters by liquid chromatography and its application to the study of the stereospecificities of D- and L-bifunctional proteins in bile acid biosynthesis
T. Kurosawa, M. Sato, K. Inoue, T. Yoshimura, M. Tohma (Hokkaido, Japan), L. Ling Jiang and T. Hashimoto (Nagano, Japan) 249

Novel instrumentation for determination of ethanol concentrations in human perspiration by gas chromatography and a good interrelationship between ethanol concentrations in sweat and blood
T. Kamei (Nagoya, Japan), T. Tsuda, Y. Mibu, S. Kitagawa, H. Wada (Nagoya, Japan), K. Naitoh and K. Nakashima (Nagoya, Japan) 259

Fine separation of protein isoforms with alternative development in two-dimensional electrophoresis
K. Nakamura and Y. Sugawara (Hiroshima, Japan) 267

Other topics

Screening of single *Escherichia coli* in a microchannel system by electric field and laser tweezers
K. Morishima, F. Arai, T. Fukuda, H. Matsuura and K. Yoshikawa (Nagoya, Japan) 273

Electrical potential oscillation across a water/octanol/water liquid membrane in the presence of two taste substances
K. Arai, F. Kusu and K. Takamura (Tokyo, Japan) 279

Clinical analysis of aldose reductase for differential diagnosis of the pathogenesis of diabetic complication
T. Tanimoto, K. Maekawa, S. Okada (Osaka, Japan) and C. Yabe-Nishimura (Kyoto, Japan) 285

Sedative effect on humans of inhalation of essential oil of linalool: Sensory evaluation and physiological measurements using optically active linalools
Y. Sugawara, C. Hara, K. Tamura, T. Fujii, K. Nakamura, T. Masujima and T. Aoki (Hiroshima, Japan) 293

Kinetic study of methionine oxidation in human parathyroid hormone
Y. Nabuchi, E. Fujiwara, H. Kuboniwa, Y. Asoh and H. Ushio (Shizuoka, Japan) 301

Minimization of genomic human hepatitis delta virus ribozyme
H. Fauzi (Tsukuba Science City, Japan and Depok, Indonesia), A. Chiba (Tsukuba Science City, Japan and Ami, Japan), F. Nishikawa, M. Roy (Tsukuba Science City, Japan), J. Kawakami (Kobe, Japan) and S. Nishikawa (Tsukuba Science City, Japan) 309

Physical stimuli and emotional stress-induced sweat secretions in the human palm and forehead
T. Kamei, T. Tsuda, S. Kitagawa, K. Naitoh, K. Nakashima (Nagoya, Japan) and T. Ohhashi (Matsumoto, Japan) 319

Author Index 327